



IMPACT OF PARTHENIUM HYSTEROPHORUS L. ON NATIVE SOIL ALGAL FLORA

D.S. Suryawanshi

Department of Botany, Jawahar college, Andur Tq. Tuljapur Dist. Osmanabad(MS).

ABSTRACT :

Soil was amended with powder from air-dried leaves, inflorescences or roots of *Parthenium hysterophorus* L. at 1, 2 and 5 per cent (w/w) levels, and inhibitory effect towards indigenous algal population was assessed. The inhibition of algae was more pronounced in the soil amended with leaf material followed by treatments with inflorescence and least with root powder. Higher levels (2 and 5%) of leaf and inflorescence amendment resulted in a greater inhibition of the algae.

KEYWORDS : amendment, inhibition, *Parthenium hysterophorus* L., soil algae.

INTRODUCTION

Parthenium hysterophorus L., a noxious weed, occurring widely in tropics and sub-tropics, is a major problem in India and Australia (Khosla and Sobti, 1979; McFadyan, 1984). Allelopathic activity of different parts of this cosmopolitan weed has been well documented (Mall and Dagar, 1979; Rajan, 1973; Sarma et al., 1976; Sukhada and Jayachandra, 1980). In pure culture studies, inhibition of growth of *Rhizobium phaseoli* and *Azotobacter vinelandii*, and activity of *Nitrosomonas* (Sukhada and Jayachandra, 1981) and antifungal properties (Char and Bhat, 1975) of *Parthenium* were also reported. Studies on the inhibitory effects of seed plants on soil algae, an ecologically beneficial group of the soil ecosystem, are very limited (Parks and Rice, 1969; Rice, 1974). There are no reports on the toxic effects of the residues of *Parthenium* on soil algae. The intent of the present study is, therefore, to evaluate the possible toxic effect of different parts of the weed on soil algal population under laboratory conditions.


MATERIALS AND METHODS

A black soil, collected from a fallow cotton field to a depth of 8cm, was air-dried and passed through a 2 mm mesh sieve. The soil had pH 7.4 (1:1.25 soil to water ratio), 1.82% organic matter and 0.044% total nitrogen.

Plant samples of *Parthenium hysterophorus* were collected at the time of flowering. The leaves, inflorescence and roots were separated, air-dried, ground to a fine powder and sieved.

From the above soil sample, 20-g portions were taken in tubes (25 x 150mm), amended with either the leaf or inflorescence or root powder to provide final concentrations of 1, 2 and 5% (w/w) on soil dry weight basis. Each portion was thoroughly mixed to ensure homogeneity. Unamended soil samples served as controls. In all the treatments and controls the soil moisture potential was maintained at 0.075 M Pa.

The tubes were incubated at room temperature ($28 \pm 4^\circ\text{C}$) and samples were withdrawn after 15 and 30 days of incubation for estimation and characterization of algae by the most probable number (MPN) method, as outlined by Muralikrishna and Venkateswarlu (1984). Fiducial limits for the estimates of viable counts at the



95% confidence level were also calculated following the formula of Fisher and Yates (1963). Most of the predominant algal forms were identified, at least to the generic level.

RESULT

The viable counts of algae in the soil were affected differentially under leaf, inflorescence and root-amendments of *Parthenium hysterophorus* (Table 1). In general, the inhibitory effect was in the order: leaf > inflorescence > root. Even a 1% amendment significantly inhibited the algae in soil. Increased toxicity towards the algae was observed with increasing levels of each plant part. The exact MPN could not be calculated with 2 and 5% amendments of leaf and inflorescence after 15 days of incubation as only less than 8 tubes were scored positive for algae out of 30 MPN tubes employed. Leaf-amendment, at the 5% level, resulted in total elimination of algae by 30 days after treatment and only one MPN tube was scored positive for algae in amendments with inflorescence at the end of 30 days as against six positive tubes by the end of 15 days. Root-amendment, at 5%, also exhibited a similar toxic effect in the 30 day sampling.

In untreated soil samples 10 genera of algae were present, 7 belonging to the Cyanophyta and 3 to the Chlorophyta (Table 2). Only the filamentous forms were encountered consistently with all the three amendments compared to the unicellular forms like *Chlorella vulgaris*, *Chlorococcum humicola*.

DISCUSSION

While demonstrating the allelopathic effects of *Parthenium hysterophorus*, Sukhada and Jayachandra (1980) reported the accumulation of large quantities of inhibitors, such as parthenin, a ses-qui-terpene lactone and phenolics in leaves followed by lower quantities in inflorescence, root and stem. The levels of toxicity caused by different plant parts reported in the present study conforms to this observation. Ravindranath (1981) opined that the inhibitors are synthesized in the leaf and then translocated to other parts of the plant.

Accumulation of leaf material at higher concentrations in the soil can be expected as a dense stand of *Parthenium* produces about 333 g m⁻² of dry leaves (Sukhada and Jayachandra, 1980). Air-dried plant parts contribute large amounts of phenols and sesquiterpene lactones to soil by rain wash or by death and decay of plant residues of *Parthenium* (Sukhada and Jayachandra, 1980).

The increased algicidal effect associated with 5%



Principal

Jawahar Arts, Science & Commerce College,
Andur Tal. Tuljapur Dist, Osmanabad

Principal

Jawahar Arts, Science & Commerce College,
Andur Tal. Tuljapur Dist, Osmanabad

Table 1. Algal population in soil as affected by the amendments of leaf, inflorescence and root of *Parthenium hysterophorus* L.

Soil amendment (per cent)	MPN (x 10 ³ g ⁻¹ soil) of algae ^a					
	15 ^b		30			
	Estimate	95% Fiducial limits		Estimate	95% Fiducial limits	
		Upper	Lower		Upper	Lower
Unamended	13.6	25.1 11.7	7.4	11.7	21.6	6.4
<i>Leaf</i>						
1	6.4	11.7	<4.1	6.4	11.7	<4.1
2	<4.2 ^c	<7.0 ^b	<4.1		<7.0	<4.1
5	<4.2	<7.0	<4.1		0	0
<i>Inflorescence</i>						
1	4.6	8.4	<4.1	<4.2	<7.0	<4.1
2	<4.2	<7.0	<4.1	<4.2	<7.0	<4.1
5	<4.2	<7.0	<4.1	<4.2	<7.0	<4.1
<i>Root</i>						
1	8.7	16.0	4.7	8.7	16.0	4.7
2	5.4	10.0	<4.1	5.4	10.0	<4.1
5	4.6	8.4	<4.1	<4.2	<7.0	<4.1

* 0-day (initial) population was 13.6 x 10³g⁻¹soil. ^b Sampling time, in days after amendment.

^c Not feasible to calculate exact MPN or fiducial limits as less than 8 tubes were scored positive.

Table 2. Occurrence of algae in soils amended with different parts of *Parthenium hysterophorus*

Organism	Unamended		Leaf						Inflorescence						Root					
	15	30	15		30		15		30		15		30		15		30			
			1	2	5	1	2	5	1	2	5	1	2	5	1	2	5	1	2	5
<i>Anabaena variabilis</i>	+	+	+	-	-	+	-	-	+	+	+	+	+	+	-	+	+	+	+	
<i>Chloralla vulgaris</i>	+	-	-	+	-	-	-	-	+	-	+	-	-	-	-	+	-	-	-	
<i>chlorococcum lmicola</i>	-	+	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	
<i>Gleocystis gigas</i>	+		+	+	-	-	-	-	-	-	+	-	-	-	+	+	-	-	-	
<i>Lyngbya gracilis</i>	+	+	-	-	-	+	-	+	-	+	-	+	+	+	-	+	+	+	+	
<i>Nostoc sp.</i>	+	+	-	-	-	+	-	-	-	-	-	-	-	-	+	+	-	-	-	
<i>Oscillatoria sp</i>	+	+	-	-	-	+	-	-	+	+	+	-	+	-	+	-	+	-	+	
<i>Phormidium tenue</i>	+	+	+	+	+	+	-	+	+	+	+	+	+	+	-	+	+	+	+	
<i>Spirulina sp</i>	-	+	-	-	1	-	-	-	-	-	-	-	-	-	+	-	-	p	-	
<i>Synchococcus elongatus</i>	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

a. Sampling time, in days, after incubation.

b. Soil amendment (per cent, s/w basis).


amendment of leaf material might be due to the release of large amounts of inhibitors into the soil after its partial or complete decomposition.

Dedonder and Van Sumere (1971) observed stimulated respiration in *Chlorella vulgaris* at higher concentrations of the phenolics which ultimately caused growth inhibition of the alga. This could be a possible reason for the complete elimination of soil algae observed here at the end of 30 days after leaf amendment. The occurrence of *Anabaena va-riabilis*, *Phormidium tenue* and *Lyngbya gracilis* even at higher levels of the plant parts indicates that filamentous blue-greens are relatively more tolerant to the inhibitors of the weed.

The present investigation reveals that inhibitors from *Parthenium hysterophorus* reaching soil through different plant parts would exert pronounced inhibitory effects on beneficial algae in soil.

REFERENCES

- Char MBS and Bhal S S 1975 Antifungal activity of pollen. *Naturwissenschaften*. 62, 536-538.
- Dedonder A and Van Sumere C F 1971 The effect of phenolics and related compounds on the growth and respiration of *Chlorella vulgaris*. *Z. Pflanzen Physiol.* 65, 70-80.
- Fisher R A and Yates F 1963 *Statistical Tables for Biological, Agricultural and Medical Research*, Oliver and Boyd, Edinburgh and London.
- Khosla S N and Sobti S N 1979 *Parthenium* — a national health hazard, its control and utility — a review. *Pesticides* 13, 121-127.
- Mall L P and Dagar J C 1979 Effect of *Parthenium hysterophorus* extract on the germination and early seedling growth of three crops. *J. Ind. Bot. Soc.* 58, 40-43.
- McFadyan R E 1984 *Parthenium* weed. *Aust. Weeds*. 3, 2-4.
- Muralikrishna P V G and Venkateswarlu K 1984 Effect of insecticides on soil algal population. *Bull. Environ. Contam. Toxicol.* 33, 241-245.
- Parks J M and Rice E L 1969 Effects of certain plants of old-field succession on the growth of blue-green algae. *Bull. Torrey Bot. Club.* 96. 345-360.
- Rajan L 1973 Growth inhibitors from *Parthenium hysterophorus*. *Curr. Sci.* 42, 729-730.
- Ravindranath 1981 Note on the effect of *Parthenium* extract on seed germination and seedling growth in crops. *Ind. J. Agric. Sci.* 51, 601-603.
- Rice E L 1974 *Allelopathy*. Academic press, INC, New York,
- Sarma K V V, Giri G S and Subrahmanyam K 1976 All-eloathic potential of *Parthenium hysterophorus* L. on seed germination and dry matter production in *Arachis hypogaea* Willd., *Crotalaria juncea* L. and *Phaseolus mungo* L. *Tropical Ecology.* 17, 76-77.
- Dhawan, S.R. AND P. Dhawan, (1996). Regeneration parthenium hysreopho L. *World Weed*, 2:244-249.
- Rivero, C.T. Chirenje, L.Q. and g. Martinez, (2004). Influence of compost on soil organic matter quality under tropical condition *Geoderma*, 123:355-361.
- Weber, J. A. Karczewska, J Drozd, M. Licznar S. Licznar and E. Jamroj, (2007). Agricultural and ecological aspects of a sandy soil as affected by the application of municipal solid waste compost. *Soil Biol. Biochem.* 39: 1294-1302 .
- Ameta S.K. Ameta, R. Soni, D. and Ameta, S.C. 2016c. Vermicomposting of *Parthenium hysterophorus* with different organic wastes and activators. *Academia Arena*, 8(4) 34-38
- Ameta S.K. Sharma S. Ameta R. and Ameta, S.C. 2015 Effect of compost of *Parthenium hysterophorus* on seed germination and survival of radish (*Raphanus sativus*): comparative study. *Int. J Bioassays.* 4(9) :4325- 4328 .
- Anabalgan, N, and Nabuvabbabm S. 2012 . Assessement impact if invasive weed *Parthenium hysterophorus* mixed with organic supplements on groth and reproduction preference of *Lampito mauritti* through vermitechnology. *Int. J. Environ. Biol.* 2(2) :88-91.



Principal

Jawahar Arts, Science & Commerce College,
Andur Tal. Tuljapur Dist, Osmanabad