

Total No. of Printed Pages: 02

SUBJECT CODE NO: - YY-2349
FACULTY OF SCIENCE AND TECHNOLOGY
B. Sc. F.Y (Sem-II)
Examination March / April - 2023
Mathematics Paper -III
Number Theory

[Time: 1:30 Hours]

[Max. Marks: 40]

Please check whether you have got the right question paper.

N. B

- 1) All questions are compulsory.
- 2) Figures to the right indicate full marks.

Q1 A. Attempt any one:

- a. If $m > 0$, then prove that $[ma, mb] = m[a, b]$ 05
- b. If p is a prime, then prove that p/a or p/b . 05

B. Attempt any one:

- c. Find the greatest common divisor of 7469 and 2464. 05
- d. Prove that $n^3 - n$ is divisible by 6. 05

Q2 A. Attempt any one:

- a. If $ax \equiv ay \pmod{m}$ and $(a, m) = 1$, then prove that $x \equiv y \pmod{m}$. 05
- b. If p is a prime, then prove that $(p-1)! \equiv -1 \pmod{p}$ 05

B. Attempt any one

- c. Prove that if p is a prime and $a^2 \equiv b^2 \pmod{p}$, then prove that $p \mid a+b$ or $p \mid a-b$. 05
- d. Find all integers that satisfy simultaneously : $x \equiv 2 \pmod{3}$,
 $x \equiv 3 \pmod{5}$, $x \equiv 5 \pmod{2}$ 05

Q3 A. Attempt any one

- a. If x is real number, then prove that $[x] \leq x < [x] + 1$, $x - 1 < [x] \leq x$, $0 \leq x - [x] < 1$. 05

- b. For every positive integer n ,
 $\sum_{d|n} \Phi(d) = n$ 05

B. Attempt any one

- c. Prove that $\mu(n)\mu(n+1)\mu(n+2)\mu(n+3) = 0$, n is positive integer. 05

- d. Find all integers x and y such that $147x + 258y = 369$ 05

Q4 Choose the correct alternative and rewrite the sentences:

10

1. If a and b are integers with $a > 0$ then there exist unique integers q and r such that $b = qa + r$, where
 - a. $0 \leq r \leq a$
 - b. $0 \leq r < a$
 - c. $0 < r \leq a$
 - d. $0 < r < a$
2. The product of any three consecutive integers is divisible by ____
 - a. 4
 - b. 5
 - c. 6
 - d. 7
3. If m is positive integer then $a \equiv b \pmod{m}$ if and only if ____
 - a. $m/a + b$
 - b. $m/a - b$
 - c. m/ab
 - d. $m/ma + b$
4. If $d(n)$ denotes the ____ positive divisors of n , then $d(12) =$ ____
 - a. 28
 - b. 24
 - c. 12
 - d. 6
5. If μ is a Mobious function then $\mu(8) =$ ____
 - a. -1
 - b. 1
 - c. 0
 - d. 8

Total No. of Printed Pages: 02

SUBJECT CODE NO: - YY-2350
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. F.Y (Sem- II)
Examination March / April - 2023
Mathematics Paper -IV Integral Calculus

[Time: 1:30 Hours]

[Max. Marks: 40]

Please check whether you have got the right question paper.

N. B

- 1) All questions are compulsory.
- 2) Figures to the right indicate full marks.

- Q1 A. Attempt any one. 05
- a) Obtain a reduction formula for $\int \cos^n x \, dx$, where n is any integer.
 - b) Evaluate : $\int \frac{1}{x^2-x-6} \, dx$.
- B. Attempt any one. 05
- c) Evaluate : $\int \frac{dx}{\sin^3 x \cos^5 x}$
 - d) Evaluate : $\int \frac{dx}{\sin x - \sin^3 x}$
- Q2 A. Attempt any one. 05
- a) Evaluate $\int_a^b x^2 \, dx$ as the limit of a sum.
 - b) Show that the intrinsic equation of the semi-cubical parabola $3ay^2 = 2x^3$ is $S = \frac{4}{9} a(\sec^3 \psi - 1)$
- B. Attempt any one. 05
- c) Find the whole length of the curve $x^{2/3} + y^{2/3} = a^{2/3}$
 - d) The part of the parabola $y^2 = 4ax$ cut by the latus rectum revolves about the tangent of the Vertex. Find the volume of the reel thus generated.
- Q3 A. Attempt any one. 05
- a) Prove that, the field \vec{F} is conservative over a region if and only if $\oint \vec{F} \cdot d\vec{r} = 0$ along any closed curve in the region.
 - b) Prove that $\int_S \vec{r} \cdot \vec{n} \, ds = 3V$, where S is closed surface, and V is volume enclosed.

B. Attempt any one:

05

c) Evaluate $\int_c \vec{F} \cdot d\vec{r}$, where $\vec{F} = x^2y^2 \hat{i} + y\hat{j}$, and the curve C is $y^2 = 4x$ in the xy-plane from (0,0) to (4,4)

d) Evaluate $\int_c \vec{F} \cdot d\vec{r}$ by stoke's theorem where $\vec{F} = y^2\hat{i} + x^2\hat{j} - (x+z)\hat{k}$ and c is the boundry of the friangle whose vertices are (0,0,0), (1,0,0) and (1,1,0)

Q4 Choose the correct alternative and rewrite the sentence.

10

1) The value of $\int_0^{\pi/2} \sin^3 x dx = \dots\dots\dots$

- a) $\frac{3}{2}$ b) $\frac{1}{3}$ c) $\frac{2}{3}$ d) $\frac{5}{3}$

2) Rectification is the process of determining

- a) The area under curve
b) The arc length of plane curve
c) Double integral
d) Multiple integral

3) $x = a(\theta - \sin\theta)$, $y = a(1 - \cos\theta)$ is an equation of

- a) Cycloid
b) Cardioid
c) Astroid
d) None of these

4) The volume of the solid generated by the revolution about the x-axis, of the area bounded by the curve $y=f(x)$, the ordinates at $x=a$, $x=b$, and the x-axis, is

- a) $\int_a^b x^2 dx$ b) $\pi \int_a^b x^2 dy$ c) $\int_a^b y dx$ d) $\pi \int_a^b y^2 dx$

5) If the circulation of a vector point function \vec{F} along any closed curve in a region is zero, then \vec{F} is said to be

- a) Irrational b) Rotational c) Solenoidal d) None of these

Total No. of Printed Pages: 2

SUBJECT CODE NO: - Y-2039
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. F.Y (Sem-I)
Examination March / April - 2023
Mathematics MAT - 101 Differential Calculus

[Time: 1:30 Hours]

[Max. Marks: 50]

Please check whether you have got the right question paper.

N. B

- 1) Attempt all questions.
- 2) Figures to the right indicate full marks.

- Q1 A) Attempt any one: 08
 a) Show that $f'(c)$, is the tangent of the angle which the tangent line to the curve $y = f(x)$ at the Point $P[c, f(c)]$ makes with x-axis.
 b) If $y = e^{ax} \sin (bx + c)$, then show that $\frac{d^n y}{dx^n} = (a^2 + b^2)^{\frac{n}{2}} e^{ax} \sin (bx + c + n \tan^{-1} (\frac{b}{a}))$
- B) Attempt any one: 07
 c) If $y = \frac{x+1}{x^2-4}$; then find $\frac{d^n y}{dx^n}$
 d) Find the value of the n^{th} derivative of $y = e^m \sin^{-1} x$ for $x = 0$.
- Q2 A) Attempt any one: 08
 a) State and prove Cauchy's mean value theorem.
 b) If $z = f(x, y)$ be a homogeneous function of x, y of degree n then prove that
 $x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = nz$,
 $\forall x, y \in$ the domain of the function.
- B) Attempt any one: 07
 c) Expand $2x^3 + 7x^2 + x - 6$ in Powers of $(x - 2)$.
 d) If $u = \tan^{-1} (\frac{x^3+y^3}{x-y})$, $x \neq y$, then show that

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = 2 \sin u \cos u$$
- Q3 A) Attempt any one: 05
 a) prove that:
 $\text{curl} (\vec{f} \times \vec{g}) = \vec{f} \text{div} \vec{g} - \vec{g} \text{div} \vec{f} + (\vec{g} \cdot \nabla) \vec{f} - (\vec{f} \cdot \nabla) \vec{g}$.
 b) Prove that:
 $\text{grad} (\phi\psi) = \phi \text{grad} \psi + \psi \text{grad} \phi$.

B) Attempt any one:

05

- c) Find $\text{grad } \log|\vec{r}|$, where $\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$.
 d) Find $\text{div } \vec{f}$ and $\text{curl } \vec{f}$, where $\vec{f} = \text{grad } (axy^2 + byz + cz^2x^3)$

Q4 Choose the correct alternative.

10

- i) If $y = x|x|$, then value of $\frac{dy}{dx}$ at the origin = _____.
 a) 1 b) x c) 0 d) 2x

- ii) If $y = \log(ax + b)$, then $\frac{d^ny}{dx^n} =$ _____.
 a) $\frac{(-1)^{n-1}(n-1)!a^n}{(ax+b)^n}$ b) $\frac{(-1)^n n! a^n}{(ax+b)^{n+1}}$ c) $\frac{(-1)^n (n-1)! a^{n-1}}{(ax+b)^n}$ d) $\frac{(-1)^{n-1} (n-1)! a^{n+1}}{(ax+b)^n}$

iii) $\log(1 + x) =$ _____.

a) $1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots + (-1)^{\frac{n}{2}} \frac{x^n}{n!} + \dots$

b) $x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots + \frac{(-1)^{n-1} x^n}{n} + \dots$

c) $1 + \frac{x^3}{3!} + \frac{x^6}{6!} + \frac{x^9}{9!} + \dots + \frac{x^{3n}}{(3n)!} + \dots$

d) $x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots + \frac{x^n}{n} + \dots$

iv) If $f(x) = |x|$, $x \in [-1, 1]$. then $f(x)$ _____

- a) Satisfy conditions of Lagrange's mean value theorem.
 b) Does not satisfy conditions of Rolle's theorem.
 c) Satisfy conditions of Rolle's theorem.
 d) Satisfy conditions of Cauchy's mean value theorem.

v) If ψ is a constant, then $\text{grad } \psi =$ _____

- a) 0 b) 1 c) -1 d) $-\psi$

Total No. of Printed Pages: 02

SUBJECT CODE NO: Y- 2040
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. F.Y (Sem-I)
Examination March / April - 2023
Mathematics MAT - 102 (Differential Equations)

[Time: 1:30 Hours]

[Max. Marks: 50]

Please check whether you have got the right question paper.

N. B

- 1) Attempt all questions.
- 2) Figures to the right indicates full marks.

Q1 A) Attempt any one. 08

a) Explain the method of solving the differential equation $\frac{dy}{dx} + py = Qy^n$, where P and Q are functions of x.

b) Explain the method of solving the differential equation.

$\frac{d^n y}{dx^n} + P_1 \frac{d^{n-1} y}{dx^{n-1}} + \dots + P_n y = X$, where P_1, P_2, \dots, P_n are constants and X is a function of x

B) Attempt any one 07

c) Solve $\frac{d^3 y}{dx^3} + \frac{d^2 y}{dx^2} - \frac{dy}{dx} - y = \cos(2x)$

d) Solve $x^2 \frac{d^2 y}{dx^2} + 7x \frac{dy}{dx} + 5y = x^5$

Q2 A) Attempt any one 08

a) Explain the method of solving the differential equation

$x^2 \frac{d^n y}{dx^n} + P_1 x^{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \dots + P_{n-1} x \frac{dy}{dx} + P_n y = X$, where P_1, P_2, \dots, P_n constants and X is a function of x.

b) Solve $\frac{d^2 y}{dx^2} - 2 \frac{dy}{dx} + y = 3e^{5/2x}$

B) Attempt any one 07

c) Solve $(2x - 1)^3 \frac{d^3 y}{dx^3} + (2x - 1) \frac{dy}{dx} - 2y = 0$

d) Solve $x^2 \frac{d^2 y}{dx^2} - x \frac{dy}{dx} + y = 2 = 2 \log x$

- Q3 A) Attempt any one 05
- a) Explain the method of solution of simultaneous differential equation

$$\frac{dx}{P} = \frac{dy}{Q} = \frac{dz}{R}$$
 where P,Q, R use functions of x,y,z
- b) With usual notation, prove that

$$\frac{1}{f(D)}(xv) = \left\{x - \frac{1}{f(D)}f'(D)\right\} \frac{1}{f(D)}V,$$
 Where V is any function of x 05
- B) Attempt any one
- c) Solve $(2ax + by + g)dx + (2cy + bx + e)dy = 0$
- d) Form the partial differential equation by eliminating the arbitrary function from
 $z = e^{ny}\phi(x - y)$
- Q4 Choose the correct alternative 10
- i. The integrating factor of differential equation $x \frac{dy}{dx} - ay = x + a$ is
- a) x^a b) $\frac{1}{x^a}$ c) $\frac{-a}{x}$ d) x^2
- ii. The partial differential equation correspond to _____
- a) Single independent variable
 b) More than one independent variable
 c) Single ordinary derivative
 d) None of these
- iii. The general solution of the differential equation
 $\frac{d^n y}{dx^n} + P_1 \frac{d^{n-1} y}{dx^{n-1}} + \dots + P_n y = X$ is
- a) $y=C.F + P.I$
 b) $y=C.F - P.I$
 c) $y=\text{complementary function}$
 d) none of these
- iv. The partial differential equation obtained by eliminating constants a and b from
 $z = a(x + y) + b$ is _____
- a) $pq=1$ b) $p=q$ c) $P^2 = q^2$ d) none of these
- v. The particular integral of the differential equation
 $\frac{d^2 y}{dx^2} + 2 \frac{dy}{dx} + y = 2e^{2x}$ is _____
- a) $\frac{2}{9}e^{2x}$ b) $\frac{1}{9}e^{2x}$ c) $2e^{2x}$ d) e^{2x}

Total No. of Printed Pages: 2

SUBJECT CODE NO: - Y-2054
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. F.Y (Sem-II)
Examination March / April - 2023
Mathematics MAT - 201 (Integral Calculas)

[Time: 1:30 Hours]

[Max. Marks: 50]

Please check whether you have got the right question paper.

N. B

- 1) Attempt all questions.
- 2) Figures to the right indicate full marks.

Q1 A) Attempt any one.

08

a) Obtain a reduction formula for $\int x^n e^{-x} dx$ and hence show that the improper integral $\int_0^{\infty} x^n e^{-x} dx = n!$ Where n is any positive integer.

b) Evaluate the definite integral

$$\int_0^{\pi/2} \sin^n x dx$$
 where n is a positive integer

B) Attempt any one.

07

c) Evaluate $\int \frac{x^5 dx}{x^3 - 2x^2 - 5x + 6}$

d) Evaluate $\int \frac{(x^3 + 2)}{(x-1)(x-2)^3} dx$

Q2 A) Attempt any one.

08

a) Evaluate $\int_a^b \cos h 2x dx$ as the limit of sum.

b) Find the area of the region lying x-axis and included between the circle $x^2 + y^2 - 2ax$ and the parabola $y^2 = ax$.

B) Attempt any one

07

c) Find the length of the arc of the curve $y = \log \tan h \left(\frac{x}{2} \right)$ from $x = 1$ to $x = 2$

d) Find the volume of the solid obtained by revolving one arc of the cycloid $x = a(\theta + \sin\theta), y = a(1 + \cos\theta)$ about $x - axis$.

Q3 A) Attempt any one

05

- a) If \vec{F} is any continuously differentiable vector point function and S is a surface bounded by curve C, then prove that $\int_C \vec{F} \cdot d\vec{r} = \int_S \text{curl } \vec{F} \cdot \vec{n} \, ds$, where the unit normal vector \vec{n} at any point of s is drawn in the sense in which a right handed screw would move when rotated in the sense of description of C.
- b) Prove that $\int_S \vec{n} \times (\vec{a} \times \vec{r}) \, ds = 2aV$.

B) Attempt any one.

05

- c) Show that $\frac{1}{3} \int_S \vec{r} \cdot d\vec{a} = V$ where V is the volume enclosed by the surface S.
- d) Evaluate $\int_C \vec{F} \cdot d\vec{r}$ where $\vec{F} = (2x^2 + y^2)\vec{i} + (3y - 4x)\vec{j}$ around the triangle ABC whose vertices are A(0,0), B(2,0), C(2,1).

Q4 Choose the correct alternatives.

10

- 1) $\int \frac{dx}{(2x-3)^3} =$
 a) $\frac{-1}{4(2x-3)^2}$ b) $\frac{1}{4(2x-3)^2}$ c) $\frac{1}{(2x-3)^2}$ d) $\frac{-1}{(2x-3)^2}$
- 2) $\int_0^{\pi/2} \sin^9 x \, dx$
 a) $\frac{315}{128}$ b) $\frac{-128}{315}$ c) $\frac{128}{315}$ d) 0
- 3) Perimeter of the cardioid $r = a(1 + \cos \theta)$ is _____
 a) 2a b) 4a c) 6a d) 8a
- 4) The volume of the solid generated by revolving the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ about the initial line is _____
 a) $\frac{\pi ab^2}{3}$ b) $3\pi ab^2$ c) πab^2 d) $\frac{4}{3}\pi ab^2$
- 5) If C be a closed curve then $\oint \vec{r} \cdot d\vec{r} =$ _____
 a) r b) r^2 c) $\frac{1}{r}$ d) 0

Total No. of Printed Pages: 03

SUBJECT CODE NO: - Y-2055
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. F.Y (Sem-II)
Examination March / April - 2023
Mathematics MAT - 202 (Geometry)

[Time: 1:30 Hours]

[Max. Marks: 50]

Please check whether you have got the right question paper.

N. B

- 1) Attempt all questions.
- 2) Figure to the right indicate full marks.

Q1 A) Attempt any one.

08

- a) Prove that every equation of the first degree in x, y, z represents a plane.
- b) Find the equations of the line passing through a given point $A(x_1, y_1, z_1)$ and having direction cosines l, m, n .

07

B) Attempt any one.

- c) Find the equation of the plane through the points $(2, 2, 1)$ and $(9, 3, 6)$ and perpendicular to the plane $2x + 6y + 6z = 9$
- d) Find two points on the line $\frac{x-2}{1} = \frac{y+3}{-2} = \frac{z-5}{2}$ on either side of $(2, -3, -5)$ and at a distance 3 from it.

Q2 A) Attempt any one.

08

- a) Find the length of the perpendicular from a given point $P(x_1, y_1, z_1)$ to a given line $\frac{x-\alpha}{l} = \frac{y-\beta}{m} = \frac{z-\gamma}{n}$

- b) Prove that a plane section of a sphere is a circle.

B) Attempt any one.

07

- c) Find the image of the point $P(1, 3, 4)$ in the plane $2x - y + z + 3 = 0$
- d) Find the equation of the sphere passing through the origin and the points $(1, 0, 0)$, $(0, 2, 0)$ and $(0, 0, 3)$

Q3 A) Attempt any one. 05

a) Find the equation of the right circular cylinder whose axis is the line

$$\frac{x-\alpha}{l} = \frac{y-\beta}{m} = \frac{z-r}{n} \text{ and whose radius is } r.$$

b) Find the points of intersection of the line $\frac{x-\alpha}{l} = \frac{y-\beta}{m} = \frac{z-r}{n}$ with the central conicoid $ax^2 + by^2 + cz^2 = 1$

B) Attempt any one.

c) Show that the distances between the parallel planes $2x - 2y + 2z + 3 = 0$ and $4x - 4y + 2z + 5 = 0$ is $\frac{1}{6}$ 05

d) Find the equation of the right circular cylinder whose radius is 2 and axis is the line $\frac{x-1}{2} = \frac{y-2}{1} = \frac{z-3}{2}$

Q4 Choose the correct alternatives and fill in the blanks. 10

1) Two planes $a_1x + b_1y + c_1z + d_1 = 0$ and $a_2x + b_2y + c_2z + d_2 = 0$ are parallel if _____.

a) $a_1 a_2 + b_1 b_2 + c_1 c_2 = 0$

b) $\frac{a_1}{a_2} + \frac{b_1}{b_2} + \frac{c_1}{c_2} = 0$

c) $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$

d) *None of these*

2) The equations to the x-axis are _____

a) $\frac{x}{0} = \frac{y}{1} = \frac{z}{1}$

b) $\frac{x}{2} = \frac{y}{0} = \frac{z}{1}$

c) $\frac{x}{1} = \frac{y}{1} = \frac{z}{0}$

d) $\frac{x}{1} = \frac{y}{0} = \frac{z}{0}$

3) The line $\frac{x+3}{3} = \frac{y-2}{-2} = \frac{z+1}{1}$ and the plane $4x + 5y + 3z - 5 = 0$ intersect at a point _____

a) (3, 1, -2)

b) (3, -2, 1)

c) (2, -1, 3)

d) (-1, -2, -3)

- 4) The radius of the sphere $x^2 + y^2 + z^2 - 2x + 4y - 6z + 7 = 0$ is _____
- a) 49
 - b) 5
 - c) -7
 - d) $\sqrt{7}$
- 5) The locus of the points of intersection of two spheres is a _____
- a) Circle
 - b) Plane
 - c) Conicoid
 - d) Cylinder

Total No. of Printed Pages: 02

SUBJECT CODE NO: - Y-2050
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. S.Y (Sem-III)
Examination March / April - 2023
Mathematics MAT - 301 Number Theory

[Time: 1:30 Hours]

[Max. Marks: 50]

Please check whether you have got the right question paper.

N. B

- i) All questions are compulsory.
 ii) Figures to the right indicate full marks.

Q1 (a) Attempt any **one** of the following: 08

- i. Given two integers a and b , with $b > 0$, then prove that there exist unique integers q and r such that

$$a = qb + r, 0 \leq r < b.$$

- ii. If $k > 0$, then prove that $gcd(ka, kb) = k gcd(a, b)$.

(b) Attempt any **one** of the following: 07

- i. By using the Euclidean algorithm, find the values of integers x and y satisfying $gcd(119, 272) = 119x + 272y$.

- ii. If a and b are both odd integers, then prove that $16 \mid a^4 + b^4 - 2$.

Q2 a) Attempt any **one** of the following: 08

- i. If $n > 1$ is a fixed integer and a, b, c, d are arbitrary integers, then prove that
 $\alphaa \equiv b \pmod{n}$ and $c \equiv d \pmod{n}$, then $a + c \equiv b + d \pmod{n}$.
 $\betaa \equiv b \pmod{n}$ and $c \equiv d \pmod{n}$, then $ac \equiv bd \pmod{n}$.

- ii. State and prove Fermat's theorem

b) Attempt any **one** of the following: 07

- i. Solve the following set of simultaneous congruences

$$x \equiv 2 \pmod{3}, x \equiv 3 \pmod{5}, x \equiv 2 \pmod{7}.$$

- ii. Find the remainder when $15!$ is divided by 17 .

- Q3 (a) Attempt any **one** of the following: 05
- Prove that the functions σ and τ are multiplicative functions.
 - If $n \geq 1$ and $\gcd(a, n) = 1$, then prove that $a^{\phi(n)} \equiv 1 \pmod{n}$.
- (b) Attempt any **one** of the following: 05
- Find $\phi(36000)$.
 - Prove that any prime of the form $3n + 1$ is also of the form $6m + 1$.
- Q4 Choose the correct alternative and **rewrite the sentence**: 10
- a) Two integers a and b not both of which are zero are said to be relatively prime, if -

- $\gcd(a, b) = 0$
 - $\gcd(a, b) = 1$
 - $\gcd(a, b) = a$
 - $\gcd(a, b) = b$
- b) If $d = \gcd(a, n)$, then the linear congruence $ax \equiv b \pmod{n}$ has a solution if and only if
- $d|a$
 - $d|b$
 - $b|d$
 - $a|d$
- c) The value of $(30) = \dots$
- 1
 - 0
 - 3
 - 1
- d) If n is even integer, then $\phi(2n) =$
- $2\phi(n)$
 - $2n$
 - n
 - $\phi(n)$
- e) A composite integer n is called a pseudoprime, if
- $n|2^n + 2$
 - $n|2^n - 2$
 - $n|2^n - 1$
 - $n|2^n$

Total No. of Printed Pages: 3

SUBJECT CODE NO: - Y-2051
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. S.Y (Sem-III)
Examination March / April - 2023
Mathematics MAT - 302 Integral Transforms

[Time:1.30 Hours]

[Max. Marks: 50]

Please check whether you have got the right question paper.

N. B

- 1) All questions are compulsory.
- 2) Figures to the right indicate full marks.

Q1 a) Attempt any one of the following: 08

- i. If
- $L^{-1}\{f(s)\} = F(t)$
- and
- $L^{-1}\{g(s)\} = G(t)$
- , then prove that

$$L^{-1}\{f(s)g(s)\} = \int_0^t F(u)G(t-u)du.$$

- ii. If
- $L\{F(t)\} = f(s)$
- , then for
- $n = 1, 2, 3, \dots$
- , prove that

$$L\{t^n F(t)\} = (-1)^n \frac{d^n}{ds^n} f(s).$$

b) Attempt any one of the following: 07

- i. If
- $L_n(x) = \frac{e^x}{n!} \frac{d^n}{dx^n} (e^{-x} x^n)$
- , then prove that
- $L\{L_n(t)\} = \frac{(s-1)^n}{s^{n+1}}$
- ,
- $s > 1$
- .

- ii. Using Heavi-side's expansion formula, find
- $L^{-1}\left\{\frac{3s+1}{(s-1)(s^2+1)}\right\}$
- .

Q2 a) Attempt any one of the following: 08

- i. If
- $F(x)$
- has the Fourier transform
- $f(s)$
- , then prove that
- $F(x) \cos ax$
- has the Fourier transform
- $\frac{1}{2}f(s-a) + \frac{1}{2}f(s+a)$
- .

- ii. If
- $L\{F(t)\} = f(s)$
- , then prove that
- $\lim_{t \rightarrow \infty} F(t) = \lim_{s \rightarrow 0} s f(s)$
- .

b) Attempt any one of the following: 07

- i. Find the Fourier transform of

$$f(x) = \begin{cases} x^2, & \text{if } |x| < a, \\ 0, & \text{if } |x| > a \end{cases}$$

- ii. Evaluate the integral
- $\int_0^\infty e^{-ax} x^{m-1} \cos bxdx$
- .

Q3 a) Attempt any one of the following: 05

i. If $L\{F(t)\} = f(s)$, then prove that $L\{e^{at}F(t)\} = f(s - a)$.

ii. If $f(s)$ is the Fourier transform of $F(x)$, then prove that $\frac{1}{s}f\left(\frac{s}{a}\right)$ is the Fourier transform of $F(ax)$.

b) Attempt any one of the following: 05

i. Find the value of $L^{-1}\left\{\frac{1}{s-2} + \frac{2}{s+5} + \frac{6}{s^4}\right\}$

ii. Prove that $L\{te^{at} \sin at\} = \frac{2a(s-a)}{(s^2-2as+2a^2)^2}$.

Q4 Choose the correct alternative and rewrite the sentence: 10

a) If $0 < l < 1$ then $\Gamma(l)\Gamma(1-l) = \dots\dots\dots$

i. $\frac{\sin l\pi}{\pi}$

ii. $\frac{\cos l\pi}{\pi}$

iii. $\frac{\pi}{\sin l\pi}$

iv. $\frac{\pi}{\cos l\pi}$

b) If $L\{F(t)\} = f(s)$ then $L\left\{\frac{F(t)}{t}\right\} = \dots\dots\dots$

i. $\int_0^\infty f(u)du$

ii. $\int_1^\infty f(u)du$

iii. $\int_s^\infty f(u)du$

iv. $\int_{-\infty}^\infty f(u)du$

c) $L^{-1}\left\{\frac{1}{s^2}\right\} = \dots\dots\dots$

i. t^2

ii. t

iii. t^3

iv. 1

d) The finite Fourier sine transform of $f(x) = 1$ for $0 < x < \pi$ is

i. $\frac{\pi(-1)^{s+1}}{s}$

ii. $\frac{\pi(-1)^{s-1}}{s}$

iii. $\frac{\pi(-1)^{s+1}}{s^2}$

iv. $\frac{1-(-1)^s}{s}$

e) $L\{\cosh at\} = \dots\dots\dots$

i. $\frac{a}{s^2-a^2}$

ii. $\frac{s}{s^2-a^2}$

iii. $\frac{a}{s^2+a^2}$

iv. $\frac{s}{s^2+a^2}$

Total No. of Printed Pages: 03

SUBJECT CODE NO: - Y-2117

FACULTY OF SCIENCE AND TECHNOLOGY

B.Sc. S.Y (Sem-III)

Examination March / April - 2023

Mathematics MAT - 303 Mechanics-I

[Time: 1:30 Hours]

[Max. Marks: 50]

Please check whether you have got the right question paper.

N. B

- 1) All questions are compulsory.
- 2) Figures to the right indicate full marks. labelled diagrams whenever necessary.

Q1 (a) Attempt any one of the following: 08

- i) Find the magnitude and direction of the resultant of any number of the coplanar forces acting at a point.
- ii) State and prove the triangle law of forces.

(b) Attempt any one of the following: 07

- i. The greatest and least magnitudes of the resultant R of two forces P and Q are G and L respectively. Show that $R^2 = G^2 \cos 2\theta + L^2 \sin 2\theta$, where 2θ is the inclination between the two forces P and Q .
- ii. A uniform plane lamina in the form of rhombus, one of whose angle is 120° , is supported by two forces applied at the centre in the direction of the diagonals so that one side of the rhombus is horizontal. Prove that if P and Q be the forces and $P > Q$ then $P = \sqrt{3}Q$.

Q2 a) Attempt any one of the following: 08

- i. Prove that the magnitude of the moment of the couple equals to the product of magnitude of a force in the couple and arm of the couple.
- ii. Prove that the sum of the vector moments of two like parallel force acting on a rigid body about any point equals to the vector moment of their resultant about the same point.

b) Attempt any one of the following: 07

- i. Three forces of magnitudes P, Q, R act along the sides BC, CA, AB of a triangle ABC, taken in order; prove that if the resultant passes through the circumcenter of AABC, then

$$P \cos A + Q \cos B + R \cos C = 0$$

- iii. Three rods of unequal lengths are joined to form a AABC. If the masses of the sides a, b, c be proportional to $(b + c - a)$, $(c + a - b)$ and $(a + b - c)$. Prove that the C.G. is incentre.

Q3 (a) Attempt any one of the following: 05

- i. If a system of parallel forces of magnitudes F_1, F_2, \dots, F_n act at some given n points, then prove that the resultant of these forces act through their centre.
- ii. A system of coplanar forces acting at a point is in equilibrium if and only if the algebraic sum of the resolved parts of the given forces along any two mutually perpendicular directions must separately vanish.

(b) Attempt any one of the following: 05

- i. Find the vector moment of a force $\vec{F} = \vec{i} + 2\vec{j} + 3\vec{k}$ acting at a point (-1, 2, 3) about origin.
- ii. If a transversal cuts the lines of action of the concurrent forces P, Q and R in A, B and C respectively, R being the resultant of the two forces P and Q. Show that

$$\frac{P}{OR} + \frac{Q}{OB} = \frac{R}{OC}$$

Q4 Choose the correct alternative and rewrite the sentence: 10

- (a) Two forces are said to be like parallel forces when they act in the and their line of action do not meet at a point.
- any direction
 - opposite direction
 - same direction
 - circular direction

(b) A particle is a body which is indefinitely small in

- i. not size and not shape
- ii. only shape
- iii. only size
- iv. size and shape

(c) The centre of the gravity is

- i. dependent
- ii. independent
- iii. not unique
- iv. unique

(d) The resolved part of the force R along the direction of the unit vector e is

- i. $\vec{R} + \vec{e}$
- ii. $\vec{R} \cdot \vec{e}$
- iii. $\vec{R} - \vec{e}$
- iv. $\vec{R} \times \vec{e}$

(e) If two forces of magnitude P each acting at an angle then the magnitude R of their resultant force is given by

- i. $R = 2P \cot \frac{\theta}{2}$
- ii. $R = 2P \tan \frac{\theta}{2}$
- iii. $R = 2 \sin \frac{\theta}{2}$
- iv. $R = 2P \cos \frac{\theta}{2}$

Total No. of Printed Pages: 2

SUBJECT CODE NO: - Y-2065
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. S.Y Sem-IV
Examination March / April - 2023
Mathematics MAT - 401 Numerical Methods

[Time: [1.30 Hours]

[Max. Marks: 50]

Please check whether you have got the right question paper.

N. B

- 1) Attempt all questions
- 2) Figure to the right indicate full marks
- 3) Use of non-Programmable calculator and logarithmic table is allowed

Q1

A) Attempt any one 08

- a) Explain the bisection method for finding real roots of an equation $f(x)=0$
- b) Derive Newton's forward difference interpolation formula

B) Attempt any one 07

- c) Use the Newton-Raphson method to find a root of the equation $x^3 - 2x - 5 = 0$ which lies between 2 and 3

d) Using Newton divided difference formula find $f(x)$ as a polynomial in x . given data is

x	-1	0	3	6	7
f(x)	3	-6	39	822	1611

Q2

A) Attempt any one 08

- a) Explain the method of fitting a straight line $Y = a_0 + a_1x$
- b) Explain the method of factorization to solve the system of linear equations

B) Attempt any one 07

- c) Determine the constant a and b by the method of least squares

Such that $y = ae^{bx}$ fits the following data

x	2	4	6	8	10
y	4.077	11.084	30.128	81.897	222.62

d) Solve the following system

$$2x+y+z=10$$

$$3x+2y+3z=18$$

$$x+4y+9z=16$$

By Gaussian elimination method

Q3 A) Attempt any one

05

a) Explain Taylor's series method to solve the differential equation

$$y' = f(x, y)$$

with the initial condition $y(x_0) = y_0$

b) With the usual notations prove that

$$\mu \equiv \left[1 + \frac{1}{4} \delta^2 \right]^{\frac{1}{2}}$$

B) Attempt any one

c) Using Picard's method obtain the solution of

$$\frac{dy}{dx} - 1 = xy \text{ with } y(0) = 1$$

05

And compute $y(0.1)$ correct to four decimal places

d) Show that

$$e^x \left(u_0 + x \Delta u_0 + \frac{x^2}{2!} \Delta^2 u_0 + \dots \right) = u_0 + u_1 x + u_2 \frac{x^2}{2!} + \dots$$

Q4 Choose the correct alternative

10

i) Which of the following is transcendental equation?

- a) $x^3 - x - 1 = 0$ b) $x^3 + x + 1 = 0$
 c) $x^3 - 2x^2 + 1 = 0$ d) $x e^x + \sin x = 0$

ii) If δ is central difference operator then $\delta y_{\frac{3}{2}} = \dots$

- a) $y_1 - y_0$ b) $y_2 - y_1$ c) $y_3 - y_2$ d) $y_4 - y_3$

iii) If $y(x) = 2x^2 + x - 1$, then $\Delta^3 y(x)$ is ----

- a) 0 b) 1 c) 2 d) 3

iv) The chebyshev polynomial of degree two is ----

- a) 1 b) x c) $2x^2 - 1$ d) $2x^2 + 1$

v) If $A = \begin{bmatrix} 1 & 0 \\ 0 & -2 \end{bmatrix}$ then the characteristic polynomial is ----

- a) $\lambda^2 + \lambda - 2$ b) $\lambda^2 - \lambda + 2$ c) $\lambda^2 + 2\lambda + 1$ d) $\lambda^2 - 2\lambda - 1$

Total No. of Printed Pages: 3

SUBJECT CODE NO: - Y-2066
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. S.Y (Sem-IV)
Examination March / April - 2023
Mathematics MAT - 402 Partial Differential Equation

[Time: 1.30 Hours]

[Max. Marks: 50]

Please check whether you have got the right question paper.

N. B

- 1) All questions are compulsory
- 2) Figures to the right indicate full marks

Q1

A) Attempt any one

- a) Explain the method of solution of the partial differential equation 08

$$f(z, p, q) = 0 \text{ and solve } p^3 + q^3 = 27z$$

- b) Define Lagrange's Linear partial differential equation. Obtain subsidiary equation of Lagrange's partial differential equation 08

B) Attempt any one

- c) Solve : 07

$$(y^2 + z^2 - x^2)p - 2xyq + 2zx = 0$$

- d) Find the complete integral of 07

$$(x + y)(p + q)^2 + (x - y)(p - q)^2 = 1$$

Q2

A) Attempt any one

- a) Explain the charpit's method for solution of partial differential equation 08

$$f(x, y, z, p, q) = 0$$

- b) Explain the method of solution of $Rr + Ss + Tt + f(x, y, z, p, q) = 0$ when $S^2 - 4RT > 0$ where R,S,T are the continuous functions of x,y and possesses continuous partial derivatives of any order 08

B) Attempt any one

- c) Find the complete integral of $x_3^2 p_1^2 p_2^2 p_3^2 + p_1^2 p_2^2 - p_3^2 = 0$ by Jacobi's method 07

- d) Solve $(D^2 - DD' + D' - 1)z = \cos(x + 2y) + e^y$ 07

Q3 A) Attempt any one 05

a) With usual notations prove that $\frac{1}{F(D,D')} (e^{ax+by} V) = e^{ax+by} \frac{1}{F(D+a,D'+b)} (v)$

b) Find the general solution of $(D - m_0 D')^2 z = 0$

B) Attempt any one 05

c) Solve $\frac{\partial q}{\partial y} - \frac{1}{y} z = x$

d) Find the general solution of the equation $(D + D')z = \sin x$

Q4 Choose the correct alternative 10

1) Auxiliary equation of

$$(y^2 + z^2 - x^2)p - 2xyq + 2xz = 0 \text{ are } \dots\dots\dots$$

a) $\frac{dx}{p} = \frac{dy}{q} = \frac{dz}{R}$

b) $\frac{dx}{y^2+z^2-x^2} = \frac{dy}{-2xy} = \frac{dz}{-2xz}$

c) $\frac{dx}{y^2+z^2+x^2} = \frac{dy}{-2xy} = \frac{dz}{2xz}$

d) $\frac{dx}{y^2+z^2-x^2} = \frac{dy}{2xy} = \frac{dz}{2xz}$

2) The symbols r,s and t are denoted respectively by -----

a) $\frac{\partial z}{\partial x}, \frac{\partial^2 z}{\partial x^2}, \frac{\partial^2 z}{\partial y^2}$

b) $\frac{\partial^2 z}{\partial y^2}, \frac{\partial^2 z}{\partial x \partial y}, \frac{\partial^2 z}{\partial x^2}$

c) $\frac{\partial^2 z}{\partial x^2}, \frac{\partial^2 z}{\partial x \partial y}, \frac{\partial^2 z}{\partial y^2}$

d) None of these

3) The complete integral of $p^2 + q^2 = n^2$ is -----

a) $z = ax + \sqrt{(n^2 - a^2)} \cdot y + c$

b) $z = ax + \sqrt{(n^2 - a^2)y} + c$

c) $z = ax + \sqrt{n^2 - a^2} + c$

d) $z = \sqrt{(n^2 - a^2)} y + c$

4) Complementary function of

$$(D^2 - 2aDD' + a^2 D'^2)z = f(y + ax) \text{ is -----}$$

a) $z = \phi_1(y + ax) + \phi_2(y + ax)$

b) $z = \phi_1(y + ax) + x\phi_2(y + ax)$

c) $z = \phi_1(y + ax) - x\phi_2(y + ax)$

d) $z = \phi_1(y - ax) + x\phi_2(y - ax)$

5) Solution of $S = 2x + 2y$ is -----

a) $z = x^2y + xy^2 + F(y) + f(x)$

b) $z = xy + xy^2 + F(y) + f(x)$

c) $z = x^2y - xy^2 + F(y) + f(x)$

d) $z = x^2y + xy^2 - F(y) - f(x)$

Total No. of Printed Pages: 2

SUBJECT CODE NO: - Y-2046
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. T.Y (Sem-V)
Examination March / April - 2023
Mathematics MAT - 501
Real Analysis – I

[Time: 1:30 Hours]

[Max. Marks: 50]

Please check whether you have got the right question paper.

N. B

1. Attempt all questions.
2. All questions carry equal marks.

Q1 A) Attempt any one:

08

- a) If $f: A \rightarrow B$ and $X \subset B, Y \subset B$, then prove that $f^{-1}(X \cap Y) = f^{-1}(X) \cap f^{-1}(Y)$
- b) If $\{S_n\}_{n=1}^{\infty}$ is a sequence of nonnegative numbers and if $\lim_{n \rightarrow \infty} S_n = L$, then prove that $L \geq 0$

B) Attempt any one:

07

- c) Show that set of all integers is countable
- d) Prove that the sequence $\left\{\frac{10^7}{n}\right\}_{n=1}^{\infty}$ has limit zero.

Q2 A) Attempt any one

08

- a) Show that the sequence $\left\{\left(1 + \frac{1}{n}\right)^n\right\}_{n=1}^{\infty}$ is convergent
- b) If $\sum_{n=1}^{\infty} a_n$ Converges to A and $\sum_{n=1}^{\infty} b_n$ converges to B, then prove that $\sum_{n=1}^{\infty} (a_n + b_n)$ converges to A+B. also if $C \in \mathbb{R}$, then prove that $\sum_{n=1}^{\infty} C a_n$ converges to CA.

B) Attempt any one

07

- c) Prove that every subsequence of a Cauchy sequence is a Cauchy sequence.
- d) Prove that the series $1 + \frac{1}{3} + \frac{1}{5} + \frac{1}{7} + \dots$ is divergent.

Q3 A) Attempt any one 05

a) If $\{S_n\}_{n=1}^{\infty}$ is a sequence of real numbers and if $\limsup_{n \rightarrow \infty} S_n = L = \liminf_{n \rightarrow \infty} S_n$ where $L \in \mathbb{R}$, then prove that $\lim_{n \rightarrow \infty} S_n = L$

b) Prove that $\frac{\partial(y_1, y_2, \dots; y_n)}{\partial(x_1, x_2, \dots; x_n)} \times \frac{\partial(x_1, x_2, \dots; x_n)}{\partial(y_1, y_2, \dots; y_n)} = 1$

B) Attempt any one. 05

c) Define i) Bounded sequence ii) Convergent sequence iii) Diverges to infinity sequence iv) Diverges to minus infinity sequence v) Monotone sequence

d) Show that the function

$$u = x + y - z, v = x - y + z,$$

$$w = x^2 + y^2 + z^2 - 2yz$$

Are not independent of one another.

Q4 Choose the correct alternative: 10

1) If $f: A \rightarrow B$ and $X \subset B$ then $f^{-1}(X) = \dots\dots\dots$

- a) $\{a \in A | f(a) \in X\}$
- b) $\{a \in B | f(a) \in B\}$
- c) $\{a \in X | f^{-1}(a) \in A\}$
- d) None of these

2) What is the value of $N \in \mathbb{I}$ such that $\left| \frac{2n}{n+3} - 2 \right| < \frac{1}{7}, n \geq N$

- a) 13 b) 23 c) 32 d) 40

3) Consider the statements

- i) Every bounded sequence is convergent
 - ii) Every subsequence of a Cauchy sequence of real number is convergent
- a) Only (i) is true
 - b) Only (ii) is true
 - c) Both (i) & (ii) are true
 - d) Both (i) & (ii) are false

4) If $\sum_{n=1}^{\infty} a_n$ is a convergent series then $\lim_{n \rightarrow \infty} a_n = L$ then $L = \dots\dots\dots$

- a) $L \neq 0$ b) $L < 0$ c) $L = 0$ d) $L > 0$

5) If $u_1 = \frac{x_2 x_3}{x_1}, u_2 = \frac{x_1 x_3}{x_2}, u_3 = \frac{x_1 x_2}{x_3}$ the $J(u_1, u_2, u_3) = \dots\dots\dots$

- a) 1 b) 2 c) 4 d) 16

Total No. of Printed Pages: 03

SUBJECT CODE NO: - Y-2047
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. T.Y (Sem-V)
Examination March / April - 2023
Mathematics MAT - 502
Abstract Algebra - I

[Time: 1:30 Hours]**[Max. Marks: 50]**

Please check whether you have got the right question paper.

N. B

- 1) All questions are compulsory.
- 2) Figures to the right indicate full marks.

Q1 A) Attempt any one:

08

- a) If H is a subgroup of a group G , then for $a, b \in G$ prove that the relation $a \equiv b \pmod{H}$ is an equivalence relation.
- b) If ϕ is a homomorphism of a group G into group \bar{G} with Kernel K , then prove that K is a normal subgroup of G .

B) Attempt any one:

07

- c) If G is a group in which $(a \cdot b)^i = a^i \cdot b^i$ for three consecutive integers i , for all $a, b \in G$, show that G is abelian.
- d) Let G be a group and g is a fixed element in G . Define $\phi: G \rightarrow G$ by $\phi(x) = gxg^{-1}$. Prove that ϕ is an isomorphism of G onto G .

Q2 A) Attempt any one:

08

- a) If ϕ is a homomorphism of a ring R into ring R' with Kernel $I(\phi)$, then prove that
 - i. $I(\phi)$ is a subgroup of R under addition.
 - ii. If $a \in I(\phi)$ and $r \in R$, then both ar and ra are in $I(\phi)$
- b) Prove that if R is a commutative ring with unit element whose only ideals are (0) and R itself, then R is a field.

B) Attempt any one:

07

- c) Prove that any field is an integral domain.
- d) If U and V are ideal of R , and if
 $U + V = \{u + v / u \in U \text{ and } v \in V\}$
 Prove that $U+V$ is also an ideal.

Q3 A) Attempt any one:

05

- a) If H and K are subgroups of a group G and $O(H) > \sqrt{O(G)}$, $O(K) > \sqrt{O(G)}$, then prove that $H \cap K \neq (e)$.
- b) If R is a commutative ring with unit element 1 and R/U is quotient ring then prove that
- R/U is commutative
 - R/U has a unit element $1+U$

B) Attempt any one:

05

- c) If N and M are normal subgroups of a group G , prove that NM is also a normal subgroup of G .
- d) If R is ring with unit element 1 and ϕ is a homomorphism of R into R' prove that $\phi(1)$ is the unit element of R'

Q4 Choose the correct alternative:

10

- i. If N is normal subgroup of a group G such that $O(G)=6$ and $O(N)=3$, then $O(G/N) = \underline{\hspace{2cm}}$
- 3
 - 2
 - 9
 - 18
- ii. For any two elements a and b of a group G , if $(a \cdot b)^2 = a^2 \cdot b^2$, then G is _____
- Abelian group
 - Quaternion group
 - Quotient group
 - None of these

- iii. If G is a group and for $x \in G$, $o(x) = n$ and $x^m = e$, then _____
- a) $m=0$
 - b) m divides n
 - c) n divides m
 - d) none of these
- iv. If an integral domain D is of finite characteristic, then its characteristic is _____
- a) A composite number
 - b) A prime number
 - c) Any integer
 - d) None of these
- v. If M is a maximal ideal of a commutative ring R with unit element, then _____
- a) R/M is a field
 - b) R/M is not a field
 - c) R is a field
 - d) None of these

Total No. of Printed Pages: 02

SUBJECT CODE NO: - Y-2061
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. T.Y (Sem-VI)
Examination March / April - 2023
Mathematics MAT-601 Real Analysis-II

[Time: 1: 30 Hours]

[Max. Marks: 50]

Please check whether you have got the right question paper.

N. B

- 1) All questions are compulsory.
- 2) Figures to the right indicate full marks.

Q1 A) Prove any one: 08

- a. Prove that every open subset G of \mathbb{R}^1 can be written $G = \cup I_n$, where I_1, I_2, \dots are a finite number or a countable number of open intervals which are mutually disjoint..
- b. Let $\langle M_1, P_1 \rangle$ and $\langle M_2, P_2 \rangle$ be metric spaces, and let $f: M_1 \rightarrow M_2$. Then prove that f is continuous on M_1 if and only if $f^{-1}(F)$ is closed subset of M_1 whenever F is a closed subset of M_2 .

B) Attempt any one 07

- c. For $P = \langle x_1, y_1 \rangle$ and $Q = \langle x_2, y_2 \rangle$, define $\sigma(P, Q) = |x_1 - x_2| + |y_1 - y_2|$, show that σ is a metric for the set of ordered pairs of real numbers.
- d. Let f be the function from \mathbb{R}^2 onto \mathbb{R}^1 defined by $f(\langle x, y \rangle) = x$ ($\langle x, y \rangle \in \mathbb{R}^2$) show that f is continuous on \mathbb{R}^2 .

Q2 A) Attempt any one 08

- a. Let $\langle M_1, P_1 \rangle$ be a compact metric space if f is a continuous function from M_1 into a metric space $\langle M_2, P_2 \rangle$, then prove that f is uniformly continuous on M_1 .
- b. If f is continuous on the closed bounded interval $[a, b]$, and if

$$F(x) = \int_a^x f(t) dt \quad (a \leq x \leq b),$$

Then prove that $F'(x) = f(x) \quad (a \leq x \leq b)$ B) Attempt any one 07

- c. Prove that every finite subset of any metric space is compact.
- d. Find the Fourier series for the function $f(x) = e^x$ in $-\pi < x < \pi$

Q3 A) Attempt any one 05

- a. if A is a closed subset of the compact metric space $\langle M, P \rangle$, then prove that the metric space $\langle A, P \rangle$ is also compact.
- b. If $f \in R[a, b]$, $g \in R[a, b]$, and if $f(x) \leq g(x)$ almost everywhere ($a \leq x \leq b$) then prove that $\int_a^b f \leq \int_a^b g$

B) Attempt any one

- c. Let $f(x) = x$ ($0 \leq x \leq 1$), Let σ be the subdivision $\{0, \frac{1}{3}, \frac{2}{3}, 1\}$ of $[0,1]$
compute $L[f; \sigma]$
- d. If $0 \leq x \leq 1$ show that
$$\frac{x^2}{\sqrt{2}} \leq \frac{x^2}{\sqrt{1+x}} \leq x^2$$

Q4 Choose the correct alternative

10

- The function P defined by $p(x, y) = |x - y|$ is a metric for the set \mathbb{R} of real numbers, then the metric space $\langle \mathbb{R}, P \rangle$ is denoted by ____
a. \mathbb{R}^d b. \mathbb{R}^d c. \mathbb{R}^1 d. \mathbb{R}^∞
- Every singleton set in a discrete metric space \mathbb{R}^d is ____
a. Open b. closed c. open and closed d. none of these
- The metric space \mathbb{R}^1 is -----
a. Not complete
b. Totally bounded
c. Complete but not totally bounded
d. Complete and totally bounded
- If f is Riemann integrable function on $[a, b]$ and $a < c < b$, then ____
a. $\int_a^b f > \int_a^c f + \int_c^b f$
b. $\int_a^b f < \int_a^c f + \int_c^b f$
c. $\int_a^b f = \int_a^c f - \int_c^b f$
d. $\int_a^b f = \int_a^c f + \int_c^b f$
- When $m=n$, for $n=0, 1, 2, \dots$
$$\int_{-\pi}^{\pi} \cos mx \cos nx \, dx = \dots$$

a. 0 b. 1 c. $-\pi$ d. π

Total No. of Printed Pages: 02

SUBJECT CODE NO: - Y-2062
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. T.Y (Sem-VI)
Examination March / April - 2023
Mathematics MAT - 602 Abstract Algebra - II

[Time: 1 :30 Hours]

[Max. Marks: 50]

Please check whether you have got the right question paper.

N. B

- 1) All questions are compulsory.
- 2) Figure to the right indicate full marks.

Q1 A. Attempt any one: 08

- a. If T is homomorphism of a vector space U onto a vector space V with kernel W , then prove that V is isomorphic to U/W
- b. Prove that if v_1, v_2, \dots, v_n are in vector space V . then either they are linearly independent or some V_k is a linear combination of the preceding ones, v_1, v_2, \dots, v_{k-1}

B. Attempt any one 07

- c. Let F be the field of all real numbers and let V be set of all sequences $\{(a_1, a_2, \dots, a_n, \dots), | a_i^2 \in F\}$
 If $U = \{(a_1, a_2, \dots, a_n, \dots), EV | \sum_{i=1}^{\infty} a_i^2 \text{ is finite}\}$ then prove that U is a subspace of V .
- d. If T is an isomorphism of vector space V onto vector space W , then prove that T maps a basis of V onto a basis of W .

Q2 A. Attempt any one: 08

- a. If W is subspace of finite-dimensional vector space V over F , then prove that $A(A(W)) = W$.
- b. Prove that if V is finite-dimensional inner product space, then V has an orthonormal set as a basis.

B. Attempt any one: 07

- c. Let V be the set of all continuous complex-valued function on the closed unit interval $[0,1]$. If $f(t), g(t) \in V$, such that

$$(f(t), g(t)) = \int_0^1 f(t) \overline{g(t)} dt$$

Prove that this define an inner product on V .

- d. If A and B are submodules of on R Modules M , then prove that $A + B = \{a + b | a \in A, b \in A\}$ is a submodule of M .

- Q3 A. Attempt any one: 05
- If W is a subspace of an inner product space V , then prove that W^\perp is a subspace of V .
 - If V is vector space over F and $v_1, v_2, \dots, v_n \in V$ are linearly independent then prove that every element in their linear span has a unique representation in the form $\lambda_1 v_1 + \lambda_2 v_2 + \dots + \lambda_n v_n$ with the $\lambda_i \in F$.
- B. Attempt any one: 05
- If V is finite-dimensional and W_1 and W_2 are subspaces of V , describe $A(W_1 \cap W_2)$ in terms of $A(W_1)$ and $A(W_2)$
 - If F is the field of real numbers, prove that the vectors $(1, 1, 0, 0)$, $(0, 1, -1, 0)$ and $(0, 0, 0, 3)$ in $F^{(4)}$ are linearly independent over F .
- Q4 Choose correct alternatives: - 10
- In an inner product space V , the inequality $|(u, v)| \leq \|u\| \cdot \|v\|$ is called ____
 - Triangle inequality
 - Bessel's inequality
 - Schwarz inequality
 - none of these
 - If V is a finite dimensional vector space and \hat{v} is its dual space then ____
 - $\text{Dim } \hat{v} = \text{Dim } V$
 - $\text{Dim } \hat{v} > \text{Dim } V$
 - $\text{Dim } \hat{v} < \text{Dim } V$
 - none of these
 - A subset S of a vector space V over F form basis if S is linearly independent and ____
 - $L(S)=S$
 - $L(S)=V$
 - $L(S)=F$
 - none of these
 - Every subspace of a vector space V other than (0) and V is called ____
 - Improper subspace
 - proper subspace
 - dual space
 - none of these
 - Vector space is defined over a ____
 - Monoids
 - group
 - ring
 - field

Total No. of Printed Pages: 02

SUBJECT CODE NO: - Y-2122
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. T.Y (Sem-VI)
Examination March / April - 2023
Mathematics
Mathematical Statistics-II - MAT -603

[Time: 1:30 Hours]

[Max. Marks: 50]

Please check whether you have got the right question paper.

N. B

- 1) All questions are compulsory.
- 2) Figures to the right indicate full marks.

Q1 A) Attempt any one:

08

a) Prove that:

The mathematical expectation of the product of a number of independent random variables is equal to the product of their expectations.

b) If x_1, x_2, \dots, x_n be n random variables, then show that

$$V\left(\sum_{i=1}^n a_i x_i\right) = \sum_{i=1}^n a_i^2 V(x_i) + 2 \sum_{i=1}^n \sum_{j=1}^n a_i a_j \text{cov}(x_i, x_j)$$

B) Attempt any one:

07

c) If m things are distributed among 'a' men and 'b' women, show that the probability that the number of things received by men is odd is given by

$$\frac{1}{2} \left[\frac{(b+a)^m - (b-a)^m}{(b+a)^m} \right]$$

d) If x be a random variable with the following probability distribution:

X: -3 6 9

P(x=x): $\frac{1}{6}$ $\frac{1}{2}$ $\frac{1}{3}$

Find E(x) and E(x²) and using the laws of expectation evaluate E(2x + 1)²

Q2 A) Attempt any one:

08

a) Find the mode of the normal distribution.

b) In case of uniform distribution, Prove that : $\mu_2 = \frac{1}{12}(b-a)^2$

B) Attempt any one:

07

c) If $x \sim B(n, p)$, show that:

$$E\left(\frac{x}{n} - p\right)^2 = \frac{pq}{n}; Cov\left(\frac{x}{n}, \frac{n-x}{n}\right) = \frac{-pq}{n}$$

- d) If x and y are independent Poisson variates with means λ_1 and λ_2 respectively
 Find i) $x + y = k$ ii) $x = y$

- Q3 A) Attempt any one 05
 a) Prove that correlation coefficient is the geometric mean between the regression coefficients.

- b) Find the mean and variance of the Poisson distribution

- B) Attempt any one: 05

- c) If x and y are independent poisson variates having mean 1 and 3 respectively.
 Find the variance of $3x + y$.

- d) If the independent random variables x ,y are binomially distributed , respectively
 $n = 3 , P = \frac{1}{3}$ and $n = 5 , P = \frac{1}{3}$, write down the probability that $x + y \geq 1$

- Q4 Choose the correct alternatives: 10

- 1) If x is a random variable , also a and b are constants , then $V(ax + b) = \dots$
 a) $a^2 V(x)$ b) $av(x) + b$ c) $V(a^2x) + b$ d) None of these

- 2) If x and y are independent the cov (x,y) =
 a) 1 b) 0 c) -1 d) 2

- 3) When the correlation coefficient $r = \pm 1$ then the two regression lines
 a) Are perpendicular to each other
 b) Coincide
 c) Are parallel to each other
 d) Do not exist

- 4) If $x \sim p(\lambda)$ then mean of poisson distribution is
 a) λ^2 b) $1/\lambda$ c) λ d) $1/\sqrt{\lambda}$

- 5) The mean of the binomial distribution is
 a) Np b) npq c) $npq(q-p)$ d) $npq\{1 + 3(n-2)pq\}$

Total No. of Printed Pages: 03

SUBJECT CODE NO: - Y-2123
FACULTY OF SCIENCE AND TECHNOLOGY
B.Sc. T.Y (Sem-VI)
Examination March / April - 2023
Ordinary Differential Equation-II - MAT- 604

[Time: 1:30 Hours]

[Max. Marks: 50]

Please check whether you have got the right question paper.

N. B

- 1) All questions are compulsory.
- 2) Figures to the right indicate full marks.

Q1 A) Attempt any one:

08

- a) Let
- $\phi_1, \phi_2, \dots, \phi_n$
- be the n solutions of

$$L(y) = y^n + a_1 y^{(n-1)} + \dots + a_n(x)y = 0 \text{ on I satisfying}$$

$$\phi_i^{(i-1)}(x_0) = 1, \phi_i^{(j-1)}(x_0) = 0, j \neq i$$

Prove that ϕ is any solution of $L(y)=0$ on I, there are n constant C_1, C_2, \dots, C_n

$$\text{Such that } \phi = C_1 \phi_1 + C_2 \phi_2 + \dots + C_n \phi_n$$

- b) Let
- $\phi_1, \phi_2, \dots, \phi_n$
- be n solutions of

$$L(y) = y^{(n)} + a_1(x)y^{(n-1)} + \dots + a_n(x)y = 0 \text{ on Interval I, and let } x_0 \text{ be}$$

$$\text{any point in I then Prove that } W(\phi_1, \phi_2, \dots, \phi_n)(x) = \exp\left[-\int_{x_0}^x a_1(t)dt\right]$$

$$W(\phi_1, \phi_2, \dots, \phi_n)(x_0)$$

B) Attempt any one

07

- c) Consider the equation

$$y'' + \frac{1}{x} y' + \frac{1}{x^2} y = 0 \text{ for } x > 0$$

I. Show that there is a solution of the form x^r , where r is constant.II. Find two linearly independent solutions for $x > 0$ and prove that they are linearly independent.III. Find two solutions ϕ_1, ϕ_2 satisfying

$$\phi_1(1) = 1, \phi_2(1) = 0$$

$$\phi_1'(1) = 0, \phi_2'(1) = 1$$

- d) Find all solutions of

$$xy'' - (x+1)y' + y = 0 \text{ given that one solution is } \phi_1(x) = e^x (x > 0)$$

Q2 A) Attempt any one

08

- a) Let b be continuous on an interval I . Let $\phi_1, \phi_2, \dots, \phi_n$ be the basis for the solution of $L(y) = y^{(n)} + a_1(x)y^{(n-1)} + \dots + a_n(x)y = 0$

Prove that every solution ψ of $L(y) = b(x)$ can be written as:

$$\psi = \psi_p + C_1\phi_1 + C_2\phi_2 + \dots + C_n\phi_n$$

Where ψ_p is particular solution of $L(y) = b(x)$ and $C_1, C_2 \dots C_n$ are constants.

Every such ψ is solution of $L(y) = b(x)$

A particular solution $\psi(P)$ is given by

$$\psi_p = \sum_{k=1}^n \phi_k(x) \int_{x_0}^x \frac{W_k(+)b(+)}{W(\phi_1, \phi_2 \dots \phi_n)t} dt$$

- b) Consider the second order Euler equation $x^2y'' + axy' + by = 0$ (a, b constant) and polynomial q is given by $q(r) = r(r-1) + ar + b$

Prove that basis for the solution of Euler equation on any interval not containing $x = 0$ is given by $\phi_1(x) = |x|^{r_1}, \phi_2(x) = |x|^{r_2}$ in case r_1 & r_2 are distinct root of q .

B) Attempt any one:

07

- c) Show that there is basis ϕ_1, ϕ_2 for the solution of

$$xy'' + 4xy' + (2 + x^2)y = 0 \quad (x > 0)$$

$$\text{of the form } \phi_1(x) = \frac{\psi_1(x)}{x^2}, \phi_2(x) = \frac{\psi_2(x)}{x^2}$$

- d) Find the linearly independent power series solution of the equation $y'' - xy = 0$

Q3 A) Attempt any one :

05

- a) Show that

$$\int_{-1}^1 P_n(x) P_m(x) dx = 0 \quad (n \neq m)$$

- b) Find all solutions of the equation $x^2y'' + 2xy' - 6y = 0 \quad (x > 0)$

B) Attempt any one

05

- c) Find the singular point of the equation $x^2y'' + (x + x^2)y' - y = 0$ and determine those which are regular singular point.

- d) Find all solutions ϕ of the form

$$\phi(x) = |x|^r \sum_{k=0}^{\infty} C_k x^k \quad (|x| > 0) \text{ for the equation}$$

$$x^2y'' + xy' + (x^2 - 1/4)y = 0$$

Q4 Choose the correct alternative

- I. If $\phi_1, \phi_2, \dots, \phi_n$ are n solutions of $L(y) = y^{(n)} + a_1(x)y^{(n-1)} + \dots + a_n(x)y = 0$ on an interval I , then they are linearly independent if and only if
- $W(\phi_1, \phi_2, \dots, \phi_n)(x) = 0 \quad \forall x \in I$
 - $W(\phi_1, \phi_2, \dots, \phi_n)(x) \neq 0 \quad \forall x \in I$
 - $W(\phi_1, \phi_2, \dots, \phi_n)(x) = \exp\left[-\int_{x_0}^x a_1(t)dt\right]$
 - $W(\phi_1, \phi_2, \dots, \phi_n)(x) = \exp\left[\int_{x_0}^x a_1(t)dt\right]$
- II. One solution of the equation $y'' - \frac{2}{x^2}y = 0$ ($0 < x < \infty$) is
- $\phi(x) = x^2$
 - $\phi(x) = x$
 - $\phi(x) = e^x$
 - $\phi(x) = e^{-x}$
- III. The singular point of the equation $a_0(x)y^n + a_1(x)y^{(n-1)} + \dots + a_n(x)y$ is the point x_0 for which
- $a_0(x_0) \neq 0$
 - $a_1(x_0) = 0$
 - $a_0(x_0) = 0$
 - $a_1(x_0) \neq 0$
- IV. $\int_{-1}^1 P_n^2(x)dx = \dots\dots\dots$
- $\frac{3}{2n+1}$
 - $\frac{1}{2n+1}$
 - $\frac{2}{2n+1}$
 - $\frac{2}{2n-1}$
- V. The equation $x^2y'' + xy' + (x^2 - a^2)y = 0$ is
- Euler equation
 - Legendre equation
 - Nonhomogeneous equation
 - Bessel equation